



Female



Male

All images Copyright Peter Drury

When Seen: Please note these dates are variable depending on the weather conditions.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
											
											
											
											

The larvae overwinters.

Food Plants:

Larvae:

	No Image	No Image	No Image
Birds Foot Trefoil (<i>Lotus corniculatus</i>)	Black Medick (<i>Medicago lupulina</i>)	Common Restharrow (<i>Ononis repens</i>)	Greater Birds-foot Trefoil (<i>Lotus pedunculatus</i>)
No Image		Intentionally Blank	Intentionally Blank
Lesser Trefoil (<i>Trifolium dubium</i>)	White Clover (<i>Trifolium repens</i>)		

Butterfly

	No Image		No Image
<u>Birds Foot Trefoil (Lotus corniculatus)</u>	<u>Bugle (Ajuga reptans)</u>	<u>Carline Thistle (Carlina vulgaris)</u>	<u>Fleabane(Pulicaria dysenterica)</u>
	(Greater Knapweed) 	No Image	No Image
<u>Kidney Vetch (Anthyllis vulneraria)</u>	<u>Knapweed (Centaurea spp.)</u>	<u>Marjoram (Origanum vulgare)</u>	<u>Ragwort (Senecio jacobaea)</u>
No Image	(Slender Thistle) 	No Image	(Common Vetch) 
<u>Thistles (Cirsium spp)</u>	<u>Thistles (Carduus spp)</u>	<u>Thyme (Thymus polytrichus)</u>	<u>Vetches (Vicia spp)</u>
	Intentionally Blank	Intentionally Blank	Intentionally Blank
<u>White Clover (Trifolium repens)</u>			

Habitat:

This species may be found in a variety of places such as grassland, waste ground, forest clearings, heathland and downland.

Most active on sunny days but roosts on dull days.

[More Common Blue butterfly images](#)

Visit my website – <http://www.imageinuk.com>