



Underside

No image

All images Copyright Peter Drury

When Seen: Please note these dates are variable depending on the weather conditions.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
											
											
											
											

Note: The pupa over winters from which the first brood emerges.

Food Plants:

Larvae

No Image		No Image	No Image
<u>Charlock (Sinapis arvensis)</u>	<u>Cuckooflower (Cardamine pratensis)</u>	<u>Garlic Mustard (Alliaria petiolata)</u>	<u>Hedge Mustard (Sisymbrium officinale)</u>
No Image	No Image	No Image	No Image
<u>Large Bitter-cress (Cardamine amara)</u>	<u>Water-cress (Rorippa nasturtium-aquaticum)</u>	<u>Wild Cabbage (Brassica oleracea)</u>	<u>Wild Radish (Raphanus raphanistrum)</u>

No Image	No Image	Intentionally Blank	Intentionally Blank
Crucifers (Cruciferae family)	Nasturtium (Tropaeolum majus)		

Butterfly

No Image	No Image		No Image
Betony (Stachys officinalis)	Bugle (Ajuga reptans)	Bluebell (Hyacinthoides non-scriptus)	Buttercups (Ranunculus spp.)
	No Image	No Image	
Cuckooflower (Cardamine pratensis)	Fleabane (Pulicaria dysenterica)	Greater Stitchwort (Stellaria holostea)	Ragged Robin (Lychnis flos-cuculi)
No Image	No Image	No Image	No Image
Ragwort (Senecio jacobaea)	Red Campion (Silene dioica)	Thistles (Cirsium spp)	Thistles (Carduus spp)
No Image	Intentionally Blank	Intentionally Blank	Intentionally Blank
Vetches (Vicia spp)			

Habitat:

This species may be seen in a variety of locations from woodland to grassland. It is particularly likes damp places (the males congregate in muddy places) but can also be seen on chalk grassland.

[More Green-veined White butterfly images](#)

Visit my website – <http://www.imageinuk.com>